arXiv:1511.06949v1 [math.CV] 22 Nov 2015

Sharp inequalities of homogeneous expansions for quasi-convex mappings of type B and almost starlike mappings of order α

Ming-Sheng Liu[1]*, Fen Wu[1] and Yan Yang[2]

1. School of Mathematical Sciences, South China Normal University, Guangzhou 510631, China

2. School of Mathematics and Computational Science, Sun Yat-Sen University, Guangzhou 510275, China

Abstract In this paper, we first obtain several sharp inequalities of homogeneous expansion for both the subclass of all normalized biholomorphic quasi-convex mappings of type B and order α and the subclass of all normalized biholomorphic almost starlike mappings of order α defined on the unit ball B of a complex Banach space X. Then, with these sharp inequalities, we derive the sharp estimates of the third and fourth homogeneous expansions for the above mappings defined on the unit polydisk D^n in $\mathbb{C}^n$.

Keywords quasi-convex mappings of type B and order α, almost starlike mappings of order α, inequalities of homogeneous expansions, the sharp estimates of the third homogeneous expansions, the sharp estimates of the fourth homogeneous expansions.

2000 Mathematics Subject Classification. 32A30, 32H02.

1 Introduction

It is well-known that there exists the following *de Branges theorem* (or *Bieberbach conjecture*) in geometric function theory of one complex variable.

Theorem A[2] If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a biholomorphic function on the unit disk $D = \{z \in \mathbb{C} : |z| < 1\}$, then

$$|a_n| \leqslant n, \quad n = 2, 3, \cdots .$$

*Corresponding author. E-mail address: liumsh@scnu.edu.com.

This research is supported by Guangdong Natural Science Foundation (Grant No.2014A030307016, 2014A030313422).

The above estimations are sharp with the Koebe function $K(z) = \frac{z}{(1-z)^2} = z + \sum_{n=2}^{\infty} n\, z^n$ as their extremal function.

Unfortunately, H. Cartan[1] has pointed out that the above theorem does not hold in the case of several complex variables. Therefore, it is necessary to require some additional properties of mappings of a class in order to obtain the analogous estimations, for instance, the starlikeness, the convexity and so on. In 1998, Sheng Gong posed the following conjecture.

Conjecture B If $f : D^n \to \mathbb{C}^n$ is a normalized biholomorphic starlike mapping, where $D^n = \{z = (z_1, \cdots, z_n) \in \mathbb{C}^n : |z_k| < 1,\ k = 1, 2, \cdots, n\}$ is the unit poly-disk in $\mathbb{C}^n$, then

$$\frac{\|D^m f(0)(z^m)\|}{m!} \leq m\|z\|^m, z \in D^n, m = 2, 3, \cdots .$$

We shall refer to Conjecture B as *Bieberbach-Gong Sheng Conjecture*. At the present time, only the case of $m = 2$ (see [3]) has been shown. However, with respect to the estimation of homogeneous expansion for normalized biholomorphic starlike mappings on the Euclidean unit ball B^n in $\mathbb{C}^n$, Roper and Suffridge in [6] have provided a counter example to verify that the above conjecture does not hold for $m = 2$. Then, in this paper, we extend some results in one complex variable to the case in several complex variables on the unit polydisk D^n in $\mathbb{C}^n$. From these results, we also obtain the sharp third estimates of homogeneous expansion for biholomorphic quasi-convex mappings of type B and order α, and the sharp third and fourth estimates of homogeneous expansion for biholomorphic almost starlike mappings of order α on the unit polydisk D^n in $\mathbb{C}^n$.

We first recall the following results in the case of one complex variable.

Theorem C[2] If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a univalent function on the unit disk D in the complex plane $\mathbb{C}$, then

$$|a_3 - a_2^2| \leq 1.$$

Theorem D[2] If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a univalent convex function on the unit disk D, then

$$|a_3 - a_2^2| \leq \frac{1 - |a_2|^2}{3}.$$

Corollary E[17] If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a univalent convex function on the unit disk D in $\mathbb{C}$, then

$$|a_3 - \frac{2}{3}a_2^2| \leq \frac{1}{3}.$$

Corollary F[17] If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a univalent starlike function on the unit disk D in $\mathbb{C}$, then

$$|2a_3 - a_2^2| \leq 2$$

Theorem G[2] If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a univalent convex function of order α on the unit disk D in $\mathbb{C}$, $0 \leq \alpha < 1$, then

$$|a_n| \leq \frac{1}{n!}\Pi_{k=2}^{n}(k - 2\alpha), n = 2, 3, \cdots .$$

The above estimation is sharp.

Theorem H[2] If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a univalent starlike function of order α on the unit disk D in $\mathbb{C}$, $0 \leq \alpha < 1$, then

$$|a_n| \leq \frac{1}{(n-1)!}\Pi_{k=2}^{n}(k - 2\alpha), n = 2, 3, \cdots .$$

The above estimation is sharp.

In the following, let X be a complex Banach space with norm $\|\cdot\|$, $B = \{x \in X : \|x\| < 1\}$ be the open unit ball in X; let $\partial_0 B$ be the boundary of B, $\bar{B}$ be the closure of B, and $\partial_0 D^n$ be the characteristic boundary (i.e. the boundary on which the maximum modulus of the holomorphic function can be attained) of the unit poly-disk D^n. $\mathbb{N}_+$ represents the set of positive integers. Let $H(B)$ be the set of all holomorphic mappings from B into X, It is well known that if $f \in H(B)$, then

$$f(y) = \sum_{n=0}^{\infty} \frac{1}{n!} D^n f(x)((y-x)^n) = \sum_{n=0}^{\infty} \frac{1}{n!} D^n f(x)(\underbrace{y-x, \cdots, y-x}_{n})$$

for all y in the neighborhood of $x \in B$, where $D^n f(x)$ is the $nth - Fréchet$ derivative of f at x. Moreover, $D^n f(x)$ is a bounded symmetric $n-$linear mapping from $\prod_{j=1}^{n} X$ into X.

A holomorphic mapping $f : B \to X$ is said to be biholomorphic if the inverse f^{-1} exists and is holomorphic on the open set $f(B)$. A mapping $f \in H(B)$ is said to be locally biholomorphic if the *Fréchet* derivative $Df(x)$ has a bounded inverse for each $x \in B$. If $f : B \to X$ is a holomorphic mapping, we say that f is normalized if $f(0) = 0$ and $Df(0) = I$, where I represents the identity operator from X into X.

If X^* is the dual space of X, for each $x \in X \backslash \{0\}$, we define

$$T(x) = \{T_x \in X^* : \|T_x\| = 1, T_x(x) = \|x\|\}.$$

According to Hahn-Banach theorem, $T(x)$ is nonempty. For any $\alpha(\neq 0) \in \mathbb{C}$, since $\frac{|\alpha|}{\alpha} T_x \in T(\alpha x)$ corresponding to each $T_x \in T(x)$, we always denote $\frac{|\alpha|}{\alpha} T_x$ by $T_{\alpha x}$.

Let $\Omega \subset \mathbb{C}^n$ be a bounded circular domain. The first order Fréchet derivative and the $m(m \geq 2)$ order Fréchet derivatives of the mapping $f \in H(\Omega)$ were denoted by $Df(z)$ and $D^m f(z)(b^{m-1}, \cdot)$, respectively. The corresponding matrixes become

$$Df(z) = \Big(\frac{\partial f_p(z)}{\partial z_k}\Big)_{1 \leq p,k \leq n},$$

$$D^m f(z)(a^{m-1}, \cdot) = \Big(\sum_{l_1.l_2,\cdots,l_{m-1}}^{n} \frac{\partial^m f_p(z)}{\partial z_k \partial z_{l_1} \cdots \partial z_{l_{m-1}}} a_{l_1} \cdots a_{l_{m-1}}\Big)_{1 \leq p,k \leq n},$$

where $f(z) = (f_1(z), f_2(z), \cdots, f_n(z))', a = (a_1, a_2, \cdots, a_n)' \in \mathbb{C}^n$.

Next, some mappings are defined as follows.

Defintion 1.1[2] Let $f : D \to \mathbb{C}$ be a holomorphic function. Then f is convex if and only if $f'(0) \neq 0$ and

$$\mathrm{Re}\Big[1 + \frac{zf''(z)}{f'(z)}\Big] \geq 0, z \in D.$$

Defintion 1.2[2] Let $f : D \to \mathbb{C}$ be a holomorphic function. We say that f is convex of order α, $0 \leq \alpha < 1$, if $f'(0) \neq 0$ and

$$\mathrm{Re}\Big[1 + \frac{zf''(z)}{f'(z)}\Big] \geq \alpha, z \in D.$$

Definition 1.3[14] Suppose that $\alpha \in [0, 1)$, and $f : B \to X$ is a locally biholomorphic mapping. If

$$\mathrm{Re}\{T_x[(Df(x))^{-1}(D^2f(x)(x^2) + Df(x)x)]\} \geq \alpha\|x\|, x \in B,$$

then f is said to be a quasi-convex mapping of type B and order α on B. Take the family as $Q_B^\alpha(B)$. When $\alpha = 0$, it is the family of quasi-convex mappings of type B.

Definition 1.4[14] Suppose that $\alpha \in [0, 1)$, and $f : B \to X$ is a locally biholomorphic mapping. If

$$\mathrm{Re}\{T_x[(Df(x))^{-1}f(x)]\} \geq \alpha\|x\|, x \in B,$$

then f is said to be an almost starlike of order α on B.

Definition 1.5[8] Suppose $f \in H(B)$. We say that $x = 0$ is the zero of order k of $f(x)$ if $f(0) = 0, \cdots, D^{k-1}f(0) = 0$, but $D^kf(0) \neq 0$, where $k \in \mathbb{N}_+$. Note that the definition is the same as that in the case $X = \mathbb{C}$.

Definition 1.6[21, 22] Suppose that $L : X^m \to \mathbb{C}$ is a continuous $m-$linear form, if

$$L(x_1, x_2, \cdots, x_m) = L(x_{\sigma(1)}, x_{\sigma(2)}, \cdots, x_{\sigma(m)}), \quad \forall x_1, \cdots, x_m \in X,$$

for any $x_1, \cdots, x_n$ in X and any permutation σ of the first m natural numbers, and

$$\|L\| = \sup\{|L(x_1, x_2, \cdots, x_m)| : \|x_1\| \leq 1, \cdots, \|x_m\| \leq 1\},$$

then L is said to be continuous symmetric $m-$linear form. Denote $\mathcal{L}^s({}^mX)$ to be the space of all continuous symmetric $m-$linear forms.

Definition 1.7[21, 22] Suppose that $L : X^m \to \mathbb{C}$ is a continuous symmetric $m-$linear form, if

$$P(x) = L(x, \cdots, x), \quad \forall x \in X,$$

then $P : X \to \mathbb{C}$ is said to be continuous homogeneous polynomial of degree m. Let $\|P\| = \sup\{|P(x)| : \|x\| \leq 1\}$. Take the family as $\mathcal{P}^s({}^mX)$.

For the sake of convenience, we let $\hat{L} = P$.

2 The inequalities of homogeneous expansion for biholomorphic quasi-convex mappings of type B and almost starlike mappings of order α

In order to prove the main results in this section, we need the following lemmas.

Lemma 2.1[2] If $f(z) = a_0 + \sum_{n=1}^{\infty} a_n z^n \in H(D)$, and $f(D) \subset D$, then

$$|a_n| \leq 1 - |a_0|^2, n = 1, 2, \cdots,$$

when $n = 1$, the above estimate is sharp.

Lemma 2.2 Let $p(z) = 1 + \sum_{n=1}^{\infty} b_n z^n \in H(D)$, and $\alpha \in [0, 1)$. If $\mathrm{Re}\, p(z) \geq \alpha, z \in D$, then

$$\Big|b_2 - \frac{b_1^2}{2(1-\alpha)}\Big| \leq 2(1-\alpha) - \frac{|b_1|^2}{2(1-\alpha)}, \tag{2.1}$$

$$\Big|b_3 - \frac{b_1 b_2}{1-\alpha} + \frac{b_1^3}{4(1-\alpha)^2}\Big| \leq 2(1-\alpha) - \frac{|b_1|^2}{2(1-\alpha)}. \tag{2.2}$$

Furthermore, we have

$$|b_2| \leq 2(1-\alpha), \tag{2.3}$$

and

$$\Big|b_2 - \frac{b_1^2}{1-\alpha}\Big| \leq 2(1-\alpha). \tag{2.4}$$

Proof Let $h(z) = \frac{1-\frac{p(z)-\alpha}{1-\alpha}}{1+\frac{p(z)-\alpha}{1-\alpha}} = \frac{1-p(z)}{1-2\alpha+p(z)}, z \in D$, then $h(0) = 0, h(D) \subset D, h(z) \in H(D)$, that is, $h(z)$ is just a $Schwarz$ function, moreover,

$$\begin{aligned} h(z) &= \frac{-\sum_{n=1}^{\infty} b_n z^n}{2(1-\alpha) + \sum_{n=1}^{\infty} b_n z^n} \\ &= \frac{-b_1}{2(1-\alpha)} z - \Big[\frac{b_2}{2(1-\alpha)} - \frac{b_1^2}{4(1-\alpha)^2}\Big] z^2 \\ &\quad - \Big[\frac{b_3}{2(1-\alpha)} - \frac{b_1 b_2}{2(1-\alpha)^2} + \frac{b_1^3}{8(1-\alpha)^3}\Big] z^3 + \cdots. \end{aligned}$$

According to Schwarz Lemma, we have

$$|h(z)/z| = 1 \ or \ |h(z)/z| < 1.$$

When $|h(z)/z| = 1$, we have

$$\Big|\frac{b_1}{2(1-\alpha)}\Big| = 1, \frac{b_2}{2(1-\alpha)} - \frac{b_1^2}{4(1-\alpha)^2} = 0, \frac{b_3}{2(1-\alpha)} - \frac{b_1 b_2}{2(1-\alpha)^2} + \frac{b_1^3}{8(1-\alpha)^3} = 0. \tag{2.5}$$

When $|h(z)/z| < 1$, we have

$$\Big|\frac{b_2}{2(1-\alpha)} - \frac{b_1^2}{4(1-\alpha)^2}\Big| \leq 1 - \Big|\frac{b_1}{2(1-\alpha)}\Big|^2, \tag{2.6}$$

$$\Big|\frac{b_3}{2(1-\alpha)} - \frac{b_1 b_2}{2(1-\alpha)^2} + \frac{b_1^3}{8(1-\alpha)^3}\Big| \leq 1 - \Big|\frac{b_1}{2(1-\alpha)}\Big|^2. \tag{2.7}$$

By (2.5), (2.6) and (2.7), we obtain

$$\Big|b_2 - \frac{b_1^2}{2(1-\alpha)}\Big| \leq 2(1-\alpha) - \frac{|b_1|^2}{2(1-\alpha)},$$
$$\Big|b_3 - \frac{b_1 b_2}{1-\alpha} + \frac{b_1^3}{4(1-\alpha)^2}\Big| \leq 2(1-\alpha) - \frac{|b_1|^2}{2(1-\alpha)}.$$

Finally, we have

$$|b_2| \leqslant \Big|b_2 - \frac{b_1^2}{2(1-\alpha)}\Big| + \frac{|b_1|^2}{2(1-\alpha)} \leqslant 2(1-\alpha),$$

and

$$\Big|b_2 - \frac{b_1^2}{1-\alpha}\Big| \leqslant \Big|b_2 - \frac{b_1^2}{2(1-\alpha)}\Big| + \frac{|b_1|^2}{2(1-\alpha)} \leq 2(1-\alpha).$$

This completes the proof.

Lemma 2.3[13] If $f(x) : B \to X$ is a normalized locally biholomorphic mapping, and $g(x) = (Df(x))^{-1}\Big(D^2 f(x)(x^2) + Df(x)x\Big)$, then

$$\frac{D^2 g(0)(x^2)}{2!} = 2\frac{D^2 f(0)(x^2)}{2!},$$
$$\frac{D^3 g(0)(x^3)}{3!} = 6\frac{D^3 f(0)(x^3)}{3!} - 4\frac{D^2 f(0)}{2!}\Big(x, \frac{D^2 f(0)(x^2)}{2!}\Big).$$

Lemma 2.4[8] If $f(x) : B \to X$ is a normalized locally biholomorphic mapping, and $g(x) = (Df(x))^{-1} f(x)$, then

$$\begin{aligned}
\frac{D^2 g(0)(x^2)}{2!} &= -\frac{D^2 f(0)(x^2)}{2!},\\
\frac{D^3 g(0)(x^3)}{3!} &= -2\frac{D^3 f(0)(x^3)}{3!} + 2\frac{D^2 f(0)}{2!}\Big(x, \frac{D^2 f(0)(x^2)}{2!}\Big),\\
\frac{D^4 g(0)(x^4)}{4!} &= -3\frac{D^4 f(0)(x^4)}{4!} + 3\frac{D^3 f(0)}{3!}\Big(x^2, \frac{D^2 f(0)(x^2)}{2!}\Big)\\
&\quad +4\frac{D^2 f(0)}{2!}\Big(x, \frac{D^3 f(0)(x^3)}{3!}\Big) - 4\frac{D^2 f(0)}{2!}\Big(x, \frac{D^2 f(0)}{2!}(x, \frac{D^2 f(0)(x^2)}{2!})\Big).
\end{aligned}$$

Now we establish the sharp inequalities of homogeneous expansion for biholomorphic quasi-convex mappings of type B and order α, and the almost starlike mappings of order α.

Theorem 2.1 If $f(x) \in Q_B^\alpha(B),\ 0 \le \alpha < 1$, then

$$\Big|T_x\Big(\frac{D^3f(0)(x^3)}{3!}\Big) - \frac{2}{3}T_x\Big(\frac{D^2f(0)}{2!}(x, \frac{D^2f(0)(x^2)}{2!})\Big)\Big| \le \frac{1-\alpha}{3}\|x\|^3, x \in B.$$

The above inequality is sharp.

Proof For fixed $x \in B\backslash\{0\}$, let $x_0 = \frac{x}{\|x\|}$. Define $p(\xi) = \frac{T_{x_0}(g(\xi x_0))}{\xi}, \xi \in D$, where

$$g(x) = (Df(x))^{-1}(D^2f(x)(x^2) + Df(x)x).$$

By the hypothesis of Theorem 2.1, we have $p(\xi) \in H(D)$, $\mathrm{Re}p(\xi) \ge \alpha$, $p(0) = 1$, and

$$p(\xi) = 1 + \frac{T_{x_0}(D^2g(0)(x_0^2))}{2!}\xi + \frac{T_{x_0}(D^3g(0)(x_0^3))}{3!}\xi^2 + \cdots .$$

According to (2.3) of Lemma 2.2, we obtain

$$\Big|T_{x_0}(\frac{D^3g(0)(x_0^3)}{3!})\Big| \le 2(1-\alpha), \quad x_0 \in \partial B.$$

From Lemma 2.3, we conclude that

$$\begin{aligned}\frac{D^3g(0)(x_0^3)}{3!} &= 6\frac{D^3f(0)(x_0^3)}{3!} - 4\frac{D^2f(0)}{2!}\Big(x_0, \frac{D^2f(0)(x_0^2)}{2!}\Big)\\ &= 6\Big[\frac{D^3f(0)(x_0^3)}{3!} - \frac{2}{3}\frac{D^2f(0)}{2!}\Big(x_0, \frac{D^2f(0)(x_0^2)}{2!}\Big)\Big].\end{aligned}$$

That is

$$\Big|T_{x_0}\Big(\frac{D^3f(0)(x_0^3)}{3!}\Big) - \frac{2}{3}T_{x_0}\Big(\frac{D^2f(0)}{2!}(x_0, \frac{D^2f(0)(x_0^2)}{2!})\Big)\Big| \le \frac{1-\alpha}{3}.$$

Hence,

$$\Big|T_x\Big(\frac{D^3f(0)(x^3)}{3!}\Big) - \frac{2}{3}T_x\Big(\frac{D^2f(0)}{2!}(x, \frac{D^2f(0)(x^2)}{2!})\Big)\Big| \le \frac{1-\alpha}{3}\|x\|^3 \ \text{ for } \ x \in B.$$

Finally, it is easy to check that the function

$$f(x) = \begin{cases} \frac{1-(1-T_u(x))^{2\alpha-1}}{2\alpha-1}x, & \alpha \in [0,1)\backslash\{\frac{1}{2}\},\\ -x\log(1-T_u(x)), & \alpha = \frac{1}{2}\end{cases}$$

satisfies the condition of Theorem 2.1, where $x \in B, u \in \partial B$. We set $x = ru, \|u\| = 1, 0 \le r < 1$. By a direct computation, we obtain that

$$\Big|T_x\Big(\frac{D^3f(0)(x^3)}{3!}\Big) - \frac{2}{3}T_x\Big(\frac{D^2f(0)}{2!}(x, \frac{D^2f(0)(x^2)}{2!})\Big)\Big| = \frac{1-\alpha}{3}r^3.$$

Hence, the inequality in the Theorem 2.1 is sharp.

Remark 2.1 Setting $X = \mathbb{C}^n, B = D^n, \alpha = 0$ in Theorem 2.1, then we can deduce Theorem 2.1 in [13].

Applying the similarly method as in the proof of Theorem 2.1, we can prove the following theorem.

Theorem 2.2 If $f(x) \in Q_B^\alpha(D^n), 0 \leq \alpha < 1$, then

$$\Big\|\frac{D^3f(0)(x^3)}{3!} - \frac{2}{3}\frac{D^2f(0)}{2!}(x, \frac{D^2f(0)(x^2)}{2!})\Big\| \leq \frac{1-\alpha}{3}\|x\|^3, \forall x \in D^n.$$

The above inequality is sharp.

Setting $n = 1$ in Theorem 2.2, we obtain Corollary 2.2.

Corollary 2.2 If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is a univalent convex function of order α on the unit disk D in the $\mathbb{C}$, $0 \leq \alpha < 1$, then

$$|a_3 - \frac{2}{3}a_2^2| \leq \frac{1-\alpha}{3}.$$

The above inequality is sharp.

Setting $\alpha = 0$ in Corollary 2.2, we obtain Corollary E.

Theorem 2.3 If $f(x)$ is an almost starlike of order α on B, $0 \leq \alpha < 1$, then

$$\begin{aligned} &\Big|2T_x\Big(\frac{D^3f(0)(x^3)}{3!}\Big)\|x\| - 2T_x\Big(\frac{D^2f(0)}{2!}(x, \frac{D^2f(0)(x^2)}{2!})\Big)\|x\| + \frac{[T_x(\frac{D^2f(0)(x^2)}{2!})]^2}{1-\alpha}\Big| \\ \leqslant\ & 2(1-\alpha)\|x\|^4, \forall x \in B. \end{aligned}$$

The above inequality is sharp.

Proof For fixed $x \in B\backslash\{0\}$, let $x_0 = \frac{x}{\|x\|}$. Define $p(\xi) = \frac{T_{x_0}(g(\xi x_0))}{\xi}, \xi \in D$, where

$$g(x) = (Df(x))^{-1}f(x).$$

By the hypothesis of Theorem 2.1, we have $p(\xi) \in H(D)$, $\mathrm{Re}p(\xi) = Re\frac{1}{|\xi|}T_{\xi x_0}(g(\xi x_0)) \geq \alpha$,

$$p(0) = \lim_{\xi\to 0}\frac{1}{\xi}T_{x_0}(g(\xi x_0)) = T_{x_0}(x_0) = 1,$$

and

$$p(\xi) = 1 + \frac{T_{x_0}(D^2g(0)(x_0^2))}{2!}\xi + \frac{T_{x_0}(D^3g(0)(x_0^3))}{3!}\xi^2 + \cdots.$$

Consequently, by (2.4) in Lemma 2.2, we obtain

$$\Big|T_{x_0}\Big(\frac{D^3g(0)(x_0^3)}{3!}\Big) - \frac{1}{1-\alpha}\Big[T_{x_0}\Big(\frac{D^2g(0)(x_0^2)}{2!}\Big)\Big]^2\Big| \leq 2(1-\alpha).$$

From Lemma 2.4, we conclude that

$$\begin{aligned} \frac{D^2g(0)(x_0^2)}{2!} &= -\frac{D^2f(0)(x_0^2)}{2!}, \\ \frac{D^3g(0)(x_0^3)}{3!} &= -2\frac{D^3f(0)(x_0^3)}{3!} + 2\frac{D^2f(0)}{2!}\Big(x_0, \frac{D^2f(0)(x_0^2)}{2!}\Big), \end{aligned}$$

that is

$$\Big|2T_{x_0}\Big(\frac{D^3f(0)(x_0^3)}{3!}\Big)-2T_{x_0}\Big(\frac{D^2f(0)}{2!}(x_0,\frac{D^2f(0)(x_0^2)}{2!})\Big)+\frac{[T_{x_0}(\frac{D^2f(0)(x_0^2)}{2!})]^2}{1-\alpha}\Big|\leq 2(1-\alpha),\forall x_0\in\partial B,$$

For $x_0=\frac{x}{\|x\|}$, so we have $T_x(\cdot)=T_{x_0}(\cdot)$. Thus we obtain

$$\Big|2T_{x}\Big(\frac{D^3f(0)}{3!}(\frac{x^3}{\|x\|^3})\Big)-2T_{x}\Big(\frac{D^2f(0)}{2!}(\frac{x}{\|x\|},\frac{D^2f(0)}{2!}(\frac{x^2}{\|x\|^2}))\Big)+\frac{[T_{x}(\frac{D^2f(0)}{2!}(\frac{x^2}{\|x\|^2}))]^2}{1-\alpha}\Big|\leq 2(1-\alpha),\forall x\in B,$$

that is,

$$\Big|2T_{x}\Big(\frac{D^3f(0)(x^3)}{3!}\Big)\|x\|-2T_{x}\Big(\frac{D^2f(0)}{2!}(x,\frac{D^2f(0)(x^2)}{2!})\Big)\|x\|+\frac{[T_{x}(\frac{D^2f(0)(x^2)}{2!})]^2}{1-\alpha}\Big|\leq 2(1-\alpha)\|x\|^4,\forall x\in B.$$

Finally, it is not difficult to check that the function

$$f(x)=\begin{cases}\frac{x}{(1-(1-2\alpha)T_u(x))^{\frac{2(1-\alpha)}{1-2\alpha}}}, & \alpha\in[0,1)\backslash\{\frac{1}{2}\},\\ xe^{T_u(x)}, & \alpha=\frac{1}{2}\end{cases}$$

satisfies the hypothesis of Theorem 2.3, where $u\in\partial B$. We set $x=ru,\|u\|=1,0\leq r<1$. By direct computation, we obtain that

$$\Big|2T_{x}\Big(\frac{D^3f(0)(x^3)}{3!}\Big)\|x\|-2T_{x}\Big(\frac{D^2f(0)}{2!}(x,\frac{D^2f(0)(x^2)}{2!})\Big)\|x\|+\frac{[T_{x}(\frac{D^2f(0)(x^2)}{2!})]^2}{1-\alpha}\Big|=2(1-\alpha)\|r\|^4.$$

Hence, the inequality in Theorem 2.3 is sharp.

Setting $\alpha=0$ in Theorem 2.3, we can obtain the following corollary.

Corollary 2.3[17] If $f(x)\in S^*(B)$. Then

$$\Big|2T_{x}\Big(\frac{D^3f(0)(x^3)}{3!}\Big)\|x\|-2T_{x}\Big(\frac{D^2f(0)}{2!}(x,\frac{D^2f(0)(x^2)}{2!})\Big)\|x\|+[T_{x}(\frac{D^2f(0)(x^2)}{2!})]^2\Big|\leq 2\|x\|^4,\forall x\in B,$$

The above inequality is sharp.

When $\alpha=0,\ n=1$, we obtain Corollary F by Corollary 2.3.

3 Applications of inequalities

In this section, with the sharp inequalities in Section 2, we will establish the sharp estimates of the third and fourth homogeneous expansions for the normalized locally biholomorphic quasi-convex mappings of type B and order α and almost starlike mappings of order α on D^n in $\mathbb{C}^n$.

In order to establish the main results in this section, we need the following lemmas.

Lemma 3.1[13] Suppose $0 \leq \alpha < 1, f : D^n \to X$ is a normalized locally biholomorphic mapping, then $f \in Q_B^\alpha(D^n)$ if and only if

$$\mathrm{Re}\Big\{\frac{g_j(z)}{z_j}\Big\} \geq \alpha, \forall z \in D^n,$$

where $g(z) = (g_1(z), g_2(z), \cdots, g_n(z))' = (Df(z))^{-1}(D^2f(z)(z^2) + Df(z)z)$ is a column vector in $\mathbb{C}^n$, $|z_j| = \|z\| = \max_{1\leq k\leq n}\{|z_k|\}$.

Lemma 3.2[13] Suppose $0 \leq \alpha < 1, f : D^n \to X$ is a normalized locally biholomorphic mapping, then f is an almost starlike mapping of order α if and only if

$$\mathrm{Re}\Big\{\frac{g_j(z)}{z_j}\Big\} \geq \alpha, \forall z \in D^n,$$

where $g(z) = (g_1(z), g_2(z), \cdots, g_n(z))' = (Df(z))^{-1}f(z))$ is a column vector in $\mathbb{C}^n$, $|z_j| = \|z\| = \max_{1\leq k\leq n}\{|z_k|\}$.

Lemma 3.3[8] Suppose $g \in H(D^n), g(0) = 0, Dg(0) = I, \alpha \in [0, 1)$. If $\mathrm{Re}\frac{g_j(z)}{z_j} \geq \alpha, z \in D^n$, where $|z_j| = \|z\| = \max_{1\leq k\leq n}\{|z_k|\}$, then

$$\frac{\|D^m g(0)(z^m)\|}{m!} \leq 2(1-\alpha)\|z\|^m, z \in D^n, m = 2, 3, \cdots.$$

The above estimate is sharp.

Lemma 3.4 If

$$\Big\|(z_1^q(\sum_{l=1}^n a_{p1l}z_l^p), z_2^q(\sum_{l=1}^n a_{p2l}z_l^p), \cdots, z_n^q(\sum_{l=1}^n a_{pnl}z_l^p))'\Big\| \leq C_0\|z\|^m,$$
$$z \in D^n, p + q = m, p, q \in \mathbb{N}^+, m = 2, 3, \cdots,$$

where each $a_{pkl}(p = 1, 2, \cdots, m-1, k, l = 1, 2, \cdots, n)$ is a complex number independent of $z_k(k = 1, 2, \cdots, n)$, $\|z\| = \max_{1\leq k\leq n}\{|z_k|\}$, C_0 is a non-negative real constant. Then

$$M = \max_{1\leq k\leq n}\{\sum_{l=1}^n |a_{pkl}|\} \leq C_0, p = 1, 2, \cdots, m-1.$$

Proof For every $z \in D^n\backslash\{0\}$, taking into the hypothesis of Lemma 3.3, we have

$$\Big|\frac{z_k^q}{\|z\|^q}\Big(\sum_{l=1}^n a_{pkl}\big(\frac{z_l}{\|z\|}\big)^p\Big)\Big| \leq C_0.$$

Especially, for each k, if $a_{pkl} \neq 0$, taking $z_l = e^{-i\frac{\arg a_{pkl}}{p}}\|z\|, l = 1, 2, \cdots, n$, where $i^2 = -1$. Then, we conclude that

$$\sum_{l=1}^n |a_{pkl}| \leq C_0, k = 1, 2, \cdots, n,$$

that is

$$M = \max_{1\le k\le n} \sum_{l=1}^{n} \{|a_{pkl}|\} \le C_0, p = 1, 2, \cdots, m-1.$$

This completes this proof.

Lemma 3.5[21] Suppose $L \in \mathcal{L}^s({}^m\ l_n^\infty)$, $\hat{L} \in \mathcal{P}^s({}^m\ l_n^\infty)$, where $\mathcal{L}^s({}^m\ l_n^\infty), \mathcal{P}^s({}^m\ l_n^\infty)$ are defined by Definitions 1.6 and 1.7, then

$$\|L\| = \|\hat{L}\|, n = 2,$$
$$\|L\| \le m^{m/2}(m+1)^{(m+1)/2}/2^m m!\|\hat{L}\|, n \ge 3, n \in \mathbb{N}_+.$$

Especially, when $m = 2$, according to Lemma 3.5, we have

Lemma 3.6 Suppose $L \in \mathcal{L}^s({}^2\ l_n^\infty)$, $\hat{L} \in \mathcal{P}^s({}^2\ l_n^\infty)$, where $\mathcal{L}^s({}^2\ l_n^\infty), \mathcal{P}^s({}^2\ l_n^\infty)$ are defined by Definitions 1.6 and 1.7, then

$$\|L\| = \|\hat{L}\|, n = 2,$$
$$\|L\| \le \frac{3}{4}\sqrt{3}\|\hat{L}\|, n \ge 3, n \in \mathbb{N}_+.$$

By Lemma 3.6, we can obtain the following lemma.

Lemma 3.7 Suppose $f : D^n \to X$ is a holomorphic mapping. Define

$$L(x, y) = D^2 f(0)(x, y), x, y \in X, \quad \|L\| = \sup\{\|L(x, y)\| : \|x\| \le 1, \|y\| \le 1\},$$
$$\hat{L}(x, x) = D^2 f(0)(x^2), x \in X, \quad \|\hat{L}\| = \sup\{\|\hat{L}(x, x)\| : \|x\| \le 1\},$$

where $\|x\| = \max_{1\le k\le n}\{|x_k|\}$, $\|y\| = \max_{1\le k\le n}\{|y_k|\}$. If L and $\hat{L}$ are bounded linear operators, then

$$\|L\| = \|\hat{L}\|, \text{ for } n = 2,$$
$$\|L\| \le \frac{3}{4}\sqrt{3}\|\hat{L}\|, \text{ for } n \ge 3,\ n \in \mathbb{N}_+.$$

Furthermore

$$\|D^2 f(0)(x, y)\| \le \|L\|\|x\|\|y\|, \text{ for } x, y \in X.$$

Proof Since L and $\hat{L}$ are bounded linear operators, according to Lemma 3.6, we have

$$\|L\| = \sup\{\|L(x, y)\| : \|x\| \le 1, \|y\| \le 1\} = \sup_{x,y\ne 0, x,y\in X} \frac{\|D^2 f(0)(x, y)\|}{\|x\|\|y\|},$$
$$\|\hat{L}\| = \sup\{\|\hat{L}(x, x)\| : \|x\| \le 1\} = \sup_{x\ne 0, x\in X} \frac{\|D^2 f(0)(x, x)\|}{\|x\|^2},$$

then

$$\|D^2f(0)(x,y)\| \le \|L\|\|x\|\|y\|, x, y \in X,$$
$$\|D^2f(0)(x,x)\| \le \|\hat{L}\|\|x\|^2, x, \in X.$$

Hence, when $\|x\| \le 1, \|y\| \le 1$, $\|D^2f(0)(x,y)\|, \|D^2f(0)(x,x)\|$ are all bounded, where

$$\|D^2f(0)(x,y)\| = \max_{1\le k\le n}\{|D^2f_k(0)(x,y)|, \|x\| \le 1, \|y\| \le 1\},$$
$$\|D^2f(0)(x,x)\| = \max_{1\le k\le n}\{|D^2f_k(0)(x^2)|, \|x\| \leqslant 1\}.$$

From Lemma 3.6, we have

$$L(x,y) = \Big(\sum_{i,j=1}^{n} a_{ij}^1 x_i y_j, \cdots, \sum_{i,j=1}^{n} a_{ij}^n x_i y_j\Big)',$$
$$\hat{L}(x,x) = \Big(\sum_{i,j=1}^{n} a_{ij}^1 x_i x_j, \cdots, \sum_{i,j=1}^{n} a_{ij}^n x_i x_j\Big)',$$

where $a_{ij}^k = \frac{\partial^2 f_k(0)}{\partial x_i \partial x_j},\ i,j,k = 1,2,\cdots,n$. Let

$$L(x,y) = (L_1(x,y), \cdots, L_n(x,y))',$$
$$\hat{L}(x,x) = (\hat{L}_1(x,x), \cdots, \hat{L_n}(x,x))',$$

where $L_k = \sum_{i,j=1}^{n} a_{ij}^k x_i y_j,\ \hat{L}_k = \sum_{i,j=1}^{n} a_{ij}^k x_i x_j,\ k = 1,2,\cdots,n$. By Definitions 1.6 and 1.7, it is easy to obtain that $L_k \in \mathcal{L}^s(^2\ l_n^\infty)$, $\hat{L}_k \in \mathcal{P}^s(^2\ l_n^\infty),\ k = 1,2,\cdots,n$. then L_k and $\hat{L}_k$ satisfy the conditions of Lemma 3.6. Therefore,

$$\|L_k\| = \|\hat{L}_k\|, \text{ for } n = 2, \tag{3.1}$$
$$\|L_k\| \le \frac{3}{4}\sqrt{3}\|\hat{L}_k\|, \text{ for } n \ge 3, n \in \mathbb{N}_+. \tag{3.2}$$

Because

$$\begin{aligned}\|L\| &= \sup_{\|x\|\le 1,\|y\|\le 1}\{\|L(x,y)\|\}\\ &= \sup_{\|x\|\le 1,\|y\|\le 1}\max_{1\le k\le n}\{|L_k(x,y)|\}\\ &\ge \sup_{\|x\|\le 1,\|y\|\le 1}\{|L_k(x,y)|\}\\ &= \|L_k\|, k = 1,2,\cdots,n,\end{aligned}$$

we have

$$\|L\| \ge \max_{1\le k\le n}\{\|L_k\|\}.$$

On the other hand,

$$\begin{aligned}
\|L\| &= \sup_{\|x\|\leq 1,\|y\|\leq 1}\{\|L(x,y)\|\} \\
&= \sup_{\|x\|\leq 1,\|y\|\leq 1}\max_{1\leq k\leq n}\{|L_k(x,y)|\} \\
&\leq \sup_{\|x\|\leq 1,\|y\|\leq 1}\max_{1\leq k\leq n}\sup_{\|x\|\leq 1,\|y\|\leq 1}\{|L_k(x,y)|\} \\
&= \sup_{\|x\|\leq 1,\|y\|\leq 1}\max_{1\leq k\leq n}\{\|L_k\|\} \\
&= \max_{1\leq k\leq n}\{\|L_k\|\}.
\end{aligned}$$

Hence,

$$\|L\| = \max_{1\leq k\leq n}\{\|L_k\|\}.$$

Similarly, we have

$$\|\hat{L}\| = \max_{1\leq k\leq n}\{\|\hat{L}_k\|\}.$$

According to (3.1) and (3.2), we obtain

$$\begin{aligned}
&\|L\| = \|\hat{L}\|, n = 2, \\
&\|L\| \leq \frac{3}{4}\sqrt{3}\|\hat{L}\|, n \geq 3, n \in \mathbb{N}_+.
\end{aligned}$$

This completes the proof.

Lemma 3.8 Assume that $\alpha \in [0, \frac{37-\sqrt{505}}{72}]$, and

$$h(x) = \frac{12\alpha^2 - 10\alpha + 1}{4(1-\alpha)^2}x^3 - \frac{1}{2(1-\alpha)}x^2 + (5-7\alpha)x + 2(1-\alpha),$$

then $h(x)$ is strictly increasing on $x \in [0, 2(1-\alpha)]$, and

$$\max_{x\in[0,2(1-\alpha)]}\{h(x)\} = h(2(1-\alpha)) = (1-\alpha)(3-4\alpha)(4-6\alpha).$$

Proof Since $h(x) = \frac{12\alpha^2-10\alpha+1}{4(1-\alpha)^2}x^3 - \frac{1}{2(1-\alpha)}x^2 + (5-7\alpha)x + 2(1-\alpha)$, we have

$$\begin{aligned}
h'(x) &= \frac{3(12\alpha^2 - 10\alpha + 1)}{4(1-\alpha)^2}x^2 - \frac{1}{1-\alpha}x + 5 - 7\alpha \\
&= \frac{3(12\alpha^2 - 10\alpha + 1)}{4(1-\alpha)^2}\Big(x - \frac{2(1-\alpha)}{3(12\alpha^2 - 10\alpha + 1)}\Big)^2 \\
&\quad +5 - 7\alpha - \frac{1}{3(12\alpha^2 - 10\alpha + 1)}.
\end{aligned}$$

When $\alpha \in [0, \frac{15-\sqrt{154}}{36})$,

$$12\alpha^2 - 10\alpha + 1 > 0, \quad 0 < \frac{2(1-\alpha)}{3(12\alpha^2 - 10\alpha + 1)} < 2(1-\alpha),$$
$$h'(\frac{2(1-\alpha)}{3(12\alpha^2 - 10\alpha + 1)}) = 5 - 7\alpha - \frac{1}{3(12\alpha^2 - 10\alpha + 1)} > 0,$$

then $h'(x) \geqslant h'(\frac{2(1-\alpha)}{3(12\alpha^2-10\alpha+1)}) > 0$ for each $x \in [0, 2(1-\alpha)]$. Hence $h(x)$ is strictly increasing on $x \in [0, 2(1-\alpha)]$.

When $\alpha \in [\frac{15-\sqrt{154}}{36}, \frac{10-\sqrt{52}}{24})$,

$$12\alpha^2 - 10\alpha + 1 > 0, \quad \frac{2(1-\alpha)}{3(12\alpha^2 - 10\alpha + 1)} \geq 2(1-\alpha), \quad h'(0) = 5 - 7\alpha > 0, \quad h'(2(1-\alpha)) > 0,$$

then $h'(x) > 0$, for every $x \in [0, 2(1-\alpha)]$. So $h(x)$ is strictly increasing on $x \in [0, 2(1-\alpha)]$.

When $\alpha = \frac{10-\sqrt{52}}{24}$,

$$h'(x) = -\frac{1}{1-\alpha}x + 5 - 7\alpha, h'(2(1-\alpha)) = 3 - 7\alpha > 0,$$

then $h'(x) > 0$, for $\forall x \in [0, 2(1-\alpha)]$, $h(x)$ is strictly increasing on $x \in [0, 2(1-\alpha)]$.

When $\alpha \in (\frac{10-\sqrt{52}}{24}, \frac{37-\sqrt{505}}{72}]$,

$$12\alpha^2 - 10\alpha + 1 < 0, \quad \frac{2(1-\alpha)}{3(12\alpha^2 - 10\alpha + 1)} < 0, \quad h'(2(1-\alpha)) \geq 0,$$

then $h'(x) > 0$, for $\forall x \in [0, 2(1-\alpha)]$, $h(x)$ is strictly increasing on $x \in [0, 2(1-\alpha)]$. Hence, for $\forall \alpha \in [0, \frac{37-\sqrt{505}}{36}]$, where $\frac{37-\sqrt{505}}{36} \approx 0.2018$, $h(x)$ is strictly increasing on $x \in [0, 2(1-\alpha)]$, and

$$\max_{x \in [0, 2(1-\alpha)]} \{h(x)\} = h(2(1-\alpha)) = (1-\alpha)(3-4\alpha)(4-6\alpha).$$

This completes this proof.

Theorem 3.1 If $0 \leq \alpha < 1$ and $f \in Q_B^\alpha(D^n)$, then for any $z \in D^n$, we have

$$\frac{\|D^3 f(0)(z^3)\|}{3!} \leq \frac{(1-\alpha)(3-2\alpha)}{3}\|z\|^3, \text{ for } n = 2, \tag{3.3}$$
$$\frac{\|D^3 f(0)(z^3)\|}{3!} \leq \frac{(1-\alpha)(\frac{3}{2}\sqrt{3} + 1 - \frac{3}{2}\sqrt{3}\alpha)}{3}\|z\|^3, \text{ for } n \geq 3, n \in \mathbb{N}_+. \tag{3.4}$$

When $n = 2$, the estimate (3.3) is sharp.

Proof For any $z \in D^n \setminus \{0\}$, define $z_0 = \frac{z}{\|z\|}$. By Lemmas 2.3 and 3.1, we have

$$\|D^2 f(0)(z_0^2)\| \leq 2(1-\alpha), \|z_0\| = 1, \tag{3.5}$$

Apparently, $\hat{L}(z^2) = D^2 f(0)(z^2)$ is a bounded linear operator. Then we have

$$\|\hat{L}\| = \sup\{\|\hat{L}(z^2)\| : \|z\| \leq 1\} = \sup\{\|\hat{L}(z^2)\| : \|z\| = 1\}. \tag{3.6}$$

According to (3.3), (3.4), we obtain

$$\|\hat{L}\| = 2(1-\alpha).$$

Let $L(x,y) = D^2 f(0)(x,y), x, y \in X$, by Lemma 3.6, we have

$$\begin{aligned} \|L\| &= 2(1-\alpha), n = 2, \\ \|L\| &\leq \frac{3}{2}\sqrt{3}(1-\alpha), n \geq 3, n \in \mathbb{N}_+ \end{aligned}$$

Let $w = \frac{D^2 f(0)(z^2)}{2!}$, again by Lemma 3.7, we obtain

$$\|D^2 f(0)(z,w)\| \leq \|L\|\|z\|\|w\| \leq 2(1-\alpha)\|z\|\|w\| \leq 2(1-\alpha)^2\|z\|^2, z \in D^n, n = 2,$$
$$\|D^2 f(0)(z,w)\| \leq \|L\|\|z\|\|w\| \leq \frac{3}{2}\sqrt{3}(1-\alpha)\|z\|\|w\| \leq \frac{3}{2}\sqrt{3}(1-\alpha)^2\|z\|^2, z \in D^n, n \geq 3, n \in \mathbb{N}_+.$$

According to Corollary 2.2, we conclude that when $n = 2$,

$$\begin{aligned} \frac{\|D^3 f(0)(z^3)\|}{3!} &\leq \frac{1-\alpha}{3}\|z\|^3 + \frac{1}{3}\|D^2 f(0)(z,w)\| \\ &\leq \frac{1-\alpha}{3}\|z\|^3 + \frac{2(1-\alpha)^2}{3}\|z\|^3 \\ &= \frac{(1-\alpha)(3-2\alpha)}{3}\|z\|^3, z \in D^2. \end{aligned}$$

Similarly, when $n \geq 3, n \in \mathbb{N}_+$,

$$\frac{\|D^3 f(0)(z^3)\|}{3!} \leq \frac{(1-\alpha)(\frac{3}{2}\sqrt{3}+1-\frac{3}{2}\sqrt{3}\alpha)}{3}\|z\|^3, z \in D^n.$$

When $n = 2$, it is easy to prove the function

$$f(z) = \begin{cases} \left(\frac{1-(1-z_1)^{2\alpha-1}}{2\alpha-1}, \frac{1-(1-z_2)^{2\alpha-1}}{2\alpha-1}\right)', & \alpha \in [0,1)\backslash\{\frac{1}{2}\}, \\ (-\log(1-z_1), -\log(1-z_2))', & \alpha = \frac{1}{2} \end{cases}$$

meets the conditions of Theorem 3.1. Taking $z = (r,0)'(0 \leq r < 1)$, then

$$\frac{\|D^3 f(0)(z^3)\|}{3!} = \frac{(1-\alpha)(3-2\alpha)}{3}r^3.$$

This completes the proof.

Theorem 3.2 Suppose $0 \leq \alpha < 1$, $f \in Q_B^\alpha(D^n)$, and $\frac{D^2 f_k(0)(z^2)}{2!} = z_k(\sum_{l=1}^n a_{kl} z_l), k = 1, 2, \cdots$, where $a_{ml} = \frac{1}{2!}\frac{\partial^2 f_m(0)}{\partial z_m \partial z_l}, m, l = 1, 2, \cdots, n$, then

$$\frac{\|D^3 f(0)(z^3)\|}{3!} \leq \frac{(1-\alpha)(3-2\alpha)}{3}\|z\|^3, z \in D^n.$$

The above estimate is sharp.

Proof. For fixed $z \in D^n \setminus \{0\}$, define $z_0 = \frac{z}{\|z\|}$. Through Lemmas 2.3, 3.1 and 3.2, we have

$$\|D^2 f(0)(z_0^2)\| \leq 2(1-\alpha).$$

According to Lemma 3.4, we obtain

$$\sum_{l=1}^{n} |a_{jl}^{(j)}| \leq 2(1-\alpha). \tag{3.7}$$

Define $w = \frac{D^2 f(0)(z^2)}{2!}$, then

$$D^2 f(0)\Big(\frac{z+w}{2}, \frac{z+w}{2}\Big) = \frac{1}{4} D^2 f(0)(z^2) + \frac{1}{2} D^2 f(0)(z,w) + \frac{1}{4} D^2 f(0)(w^2),$$

$$D^2 f(0)\Big(\frac{z-w}{2}, \frac{z-w}{2}\Big) = \frac{1}{4} D^2 f(0)(z^2) - \frac{1}{2} D^2 f(0)(z,w) + \frac{1}{4} D^2 f(0)(w^2).$$

Because $D^2 f_j(0)(z^2) = z_j\Big(\sum_{l=1}^{n} a_{jl}^{(j)} z_l\Big)$, where $|z_j| = \|z\| = \max_{1\leq k\leq n}\{|z_k|\}$, then

$$\begin{aligned} D^2 f_j(0)(z,w) &= D^2 f_j(0)\Big(\frac{z+w}{2}, \frac{z+w}{2}\Big) - D^2 f_j(0)\Big(\frac{z-w}{2}, \frac{z-w}{2}\Big) \\ &= -\frac{1}{4}(z_j - w_j)\Big(\sum_{l=1}^{n} a_{jl}^{(j)}(z_l - w_l)\Big) \\ &= \frac{1}{2}\Big[z_j\Big(\sum_{l=1}^{n} a_{jl}^{j} w_l\Big) + w_j\Big(\sum_{l=1}^{n} a_{jl}^{(j)} z_l\Big)\Big]. \end{aligned} \tag{3.8}$$

By (3.5) and (3.6), we have

$$\begin{aligned} \Big|D^2 f_j(0)\Big(z_0, \frac{D^2 f(0)(z_0^2)}{2!}\Big)\Big| &= \frac{1}{2}\Big|\frac{z_j}{\|z\|}\Big(\sum_{l=1}^{n} a_{jl}^{(j)} \frac{D^2 f_l(0)(z_0^2)}{2!}\Big) + \frac{D^2 f_j(0)(z_0^2)}{2!}\Big(\sum_{l=1}^{n} a_{jl}^{(j)} \frac{z_l}{\|z\|}\Big)\Big| \\ &\leq \frac{1}{2}\Big[\Big|\sum_{l=1}^{n} a_{jl}^{(j)} \frac{D^2 f_l(0)(z_0^2)}{2!}\Big| + \frac{|D^2 f_j(0)(z_0^2)|}{2!}\Big|\sum_{l=1}^{n} a_{jl}^{(j)} \frac{z_l}{\|z\|}\Big|\Big] \\ &\leq \frac{1}{2}\Big[\sum_{l=1}^{n} |a_{jl}^{(j)}| \frac{|D^2 f_l(0)(z_0^2)|}{2!} + \frac{|D^2 f_j(0)(z_0^2)|}{2!}\sum_{l=1}^{n} |a_{jl}^{(j)}| \frac{|z_l|}{\|z\|}\Big] \\ &\leq \frac{1}{2}\Big[2(1-\alpha)(1-\alpha) + (1-\alpha)2(1-\alpha)\Big] \\ &= 2(1-\alpha)^2, j = 1.2, \cdots, n. \end{aligned}$$

That is

$$\Big\| D^2 f(0)\Big(z_0, \frac{D^2 f(0)(z_0^2)}{2!}\Big)\Big\| \leq 2(1-\alpha)^2,\ z_0 \in \partial D^n.$$

In view of the maximum modulus theorem of holomorphic functions on the unit polydisk, we obtain

$$\Big\| D^2 f(0)\Big(z, \frac{D^2 f(0)(z^2)}{2!}\Big)\Big\| \leq 2(1-\alpha)^2\|z\|^3, z \in D^n.$$

Using Corollary 2.2, we conclude that

$$\begin{aligned}\frac{\|D^3 f(0)(z^3)\|}{3!} &\leq \frac{1-\alpha}{3}\|z\|^3 + \frac{1}{3}\Big\| D^2 f(0)\Big(z, \frac{D^2 f(0)(z^2)}{2!}\Big)\Big\| \\ &\leq \frac{1-\alpha}{3}\|z\|^3 + \frac{2(1-\alpha)^2}{3}\|z\|^3 \\ &= \frac{(1-\alpha)(3-2\alpha)}{3}\|z\|^3,\ z \in D^n.\end{aligned}$$

Finally, it is easy to prove the function

$$f(z) = \begin{cases} \Big(\frac{1-(1-z_1)^{2\alpha-1}}{2\alpha-1}, \frac{1-(1-z_2)^{2\alpha-1}}{2\alpha-1}, \cdots, \frac{1-(1-z_n)^{2\alpha-1}}{2\alpha-1}\Big)', & \alpha \in [0,1)\backslash\{\frac{1}{2}\}, \\ (-\log(1-z_1), -\log(1-z_2), \cdots, -\log(1-z_n))', & \alpha = \frac{1}{2} \end{cases}$$

meets the conditions of Theorem 3.2. Taking $z = (r, 0, \cdots, 0)'(0 \leq r < 1)$, then

$$\frac{\|D^3 f(0)(z^3)\|}{3!} = \frac{(1-\alpha)(3-2\alpha)}{3}r^3.$$

This completes the proof of Theorem 3.2.

Theorem 3.3 Suppose $0 \leq \alpha \leq \frac{1}{2}$, f is an almost starlike of order α in D^n, and

$$\begin{aligned}&2z_k\frac{D^2 f_k(0)}{2!}\Big(z_0, \frac{D^2 f(0)(z_0^2)}{2!}\Big)\|z\| - \frac{1}{1-\alpha}\Big[\frac{D^2 f_k(0)(z_0^2)}{2!}\Big]^2\|z\|^2 \\ &= \frac{1-2\alpha}{1-\alpha}z_k^2\Big(\sum_{k=1}^n a_{km}\frac{z_k}{\|z\|}(\sum_{l=1}^n a_{ml}\frac{z_l}{\|z\|})\Big),\ \text{ for } z \in D^n\backslash\{0\}.\end{aligned}$$

where $a_{ml} = \frac{1}{2!}\frac{\partial^2 f_m(0)}{\partial z_m \partial z_l}, m, l = 1, 2, \cdots, n$, then

$$\frac{\|D^3 f(0)(z^3)\|}{3!} \leq (1-\alpha)(3-4\alpha)\|z\|^3,\ \text{ for } z \in D^n.$$

The above estimate is sharp.

Proof. For fixed $z \in D^n \setminus \{0\}$, define $z_0 = \frac{z}{\|z\|}$. Taking $T_z = (0, \cdots, 0, \frac{|z_j|}{z_j}, \cdots, 0)$, where $|z_j| = \|z\| = \max_{1\leq k\leq n}\{|z_k|\}$. Applying Theorem 2.3, we have

$$\Big|2\frac{D^3 f_j(0)(z_0^3)}{3!}\frac{\|z\|}{z_j} - 2\frac{D^2 f_j(0)}{2!}\Big(z_0, \frac{D^2 f(0)(z_0^2)}{2!}\Big)\frac{\|z\|}{z_j} + \frac{1}{1-\alpha}\Big[\frac{D^2 f_j(0)(z_0^2)}{2!}\Big]^2(\frac{\|z\|}{z_j})^2\Big| \leq 2(1-\alpha).$$

According to the hypothesis of Theorem 3.3, we obtain

$$\frac{|D^3 f_j(0)(z_0^3)|}{3!} \leq 1 - \alpha + \frac{1-2\alpha}{1-\alpha}\frac{M^2}{2},$$

where $|z_j| = \|z\| = \max_{1\leq k\leq n}\{|z_k|\}$, $M = \max_{1\leq k\leq n}\{\sum_{l=1}^{n}|a_{kl}|\}$. Applying the maximum modulus principle of holomorphic functions, and Lemmas 3.1, 3.3 and 3.4, $C_0 = 2(1-\alpha)$, then

$$\frac{\|D^3f(0)(z^3)\|}{3!} \leq (1-\alpha+\frac{1-2\alpha}{1-\alpha}\frac{M^2}{2})\|z\|^3 \leq (1-\alpha)(3-4\alpha)\|z\|^3, z\in D^n.$$

Finally, it is easy to prove that the function

$$f(z) = \begin{cases} \Big(\frac{z_1}{(1-(1-2\alpha)z_1)^{\frac{2(1-\alpha)}{1-2\alpha}}}, \cdots, \frac{z_n}{(1-(1-2\alpha)z_n)^{\frac{2(1-\alpha)}{1-2\alpha}}}\Big)', & 0\leqslant\alpha<\frac{1}{2}, \\ (z_1e^{z_1}, \cdots, z_ne^{z_n})', & \alpha=\frac{1}{2} \end{cases}$$

meets the hypothesis of Theorem 3.3, where $z\in D^n$. Taking $z=(r,0,\cdots,0)'(0\leq r<1)$,

$$D^3f(0)(z^3) = (6(1-\alpha)(3-4\alpha)r^3, 0, \cdots, 0)'.$$

Then

$$\frac{\|D^3f(0)(z^3)\|}{3!} = (1-\alpha)(3-4\alpha)r^3,$$

This completes the proof of Theorem 3.3.

Corollary 3.3 Suppose $0\leq\alpha\leq\frac{1}{2}$, f is an almost starlike mapping of order α in D^n, and $\frac{D^2f_k(0)(z^2)}{2!} = z_k(\sum_{l=1}^{n}a_{kl}z_l), k=1,2,\cdots$, where $a_{ml} = \frac{1}{2!}\frac{\partial^2 f_m(0)}{\partial z_m\partial z_l}, m,l=1,2,\cdots,n$, then

$$\frac{\|D^3f(0)(z^3)\|}{3!} \leq (1-\alpha+\frac{1-2\alpha}{1-\alpha}\frac{M^2}{2})\|z\|^3 \leq (1-\alpha)(3-4\alpha)\|z\|^3, z\in D^n.$$

where $M = \max_{1\leq k\leq n}\{\sum_{l=1}^{n}|a_{kl}|\}$. The above estimate is sharp.

Remark Setting $\alpha=0$ and $f\in S^*(D^n)$ in Corollary 3.3, we can obtain Theorem 1.3 in [15].

Theorem 3.4 If $0\leq\alpha<1$, f is an almost starlike of order α in D^n, then

$$\frac{\|D^3f(0)(z^3)\|}{3!} \leq (1-\alpha)(5-4\alpha)\|z\|^3, z\in D^n, n=2,$$
$$\frac{\|D^3f(0)(z^3)\|}{3!} \leq (1-\alpha)(3\sqrt{3}+1-3\sqrt{3}\alpha)\|z\|^3, z\in D^n, n\geq 3, n\in\mathbb{N}_+.$$

Proof For fixed $z\in D^n\setminus\{0\}$, define $z_0 = \frac{z}{\|z\|}$. By Lemmas 2.3 and 3.1, we have

$$\|D^2f(0)(z_0^2)\| \leq 4(1-\alpha), \|z_0\| = 1. \tag{3.9}$$

Apparently, $\hat{L}(z^2) = D^2f(0)(z^2)$ is a bounded linear operator, we have

$$\|\hat{L}\| = \sup\{\|\hat{L}(z^2)\| : \|z\|\leq 1\} = \sup\{\|\hat{L}(z^2)\| : \|z\| = 1\}. \tag{3.10}$$

According to (3.3), (3.4), we obtain

$$\|\hat{L}\| = 4(1-\alpha).$$

Let $L(x,y)=D^2f(0)(x,y), x,y\in X$, by Lemma 3.6 we have

$$\begin{aligned}\|L\| &= 2(1-\alpha), \text{ for } n=2,\\ \|L\| &\le \frac{3}{2}\sqrt{3}(1-\alpha), \text{ for } n\ge 3, n\in\mathbb{N}_+.\end{aligned}$$

Let $w=\frac{D^2f(0)(z^2)}{2!}$ and $\|w_0\|=\frac{\|D^2f(0)(z_0^2)\|}{2!}\le 2(1-\alpha)$, again by Lemma 3.7,

$$\begin{aligned}\|D^2f(0)(z,w)\| &\le \|L\|\|z\|\|w\|\le 4(1-\alpha)\|z\|\|w\|\le 8(1-\alpha)^2\|z\|^2, z\in D^n, n=2,\\ \|D^2f(0)(z,w)\| &\le \|L\|\|z\|\|w\|\le 3\sqrt{3}(1-\alpha)\|z\|\|w\|\\ &\le 6\sqrt{3}(1-\alpha)^2\|z\|^2, z\in D^n, n\ge 3, n\in\mathbb{N}_+.\end{aligned}$$

According to Lemmas 2.2 and 2.4, we conclude that

$$\left|2\frac{D^3f(0)(z_0^3)}{3!}-2\frac{D^2f(0)}{2!}(z_0,w_0)\right|\le 2(1-\alpha). \tag{3.11}$$

By (3.7), (3.8) and (3.9), we have

$$\begin{aligned}\frac{\|D^3f(0)(z_0^3)\|}{3!}&\le 1-\alpha+\frac{1}{2}\|Df(0)(z_0,w_0)\|\le(1-\alpha)(5-4\alpha), n=2,\\ \frac{\|D^3f(0)(z_0^3)\|}{3!}&\le(1-\alpha)(3\sqrt{3}+1-3\sqrt{3}\alpha), n\ge 3, n\in\mathbb{N}_+,\end{aligned}$$

that is

$$\begin{aligned}\frac{\|D^3f(0)(z^3)\|}{3!}&\le(1-\alpha)(5-4\alpha)\|z\|^3, z\in D^n, n=2,\\ \frac{\|D^3f(0)(z^3)\|}{3!}&\le(1-\alpha)(3\sqrt{3}+1-3\sqrt{3}\alpha)\|z\|^3, z\in D^n, n\ge 3, n\in\mathbb{N}_+.\end{aligned}$$

This completes the proof.

Theorem 3.5 Suppose $0\le\alpha\le\frac{37-\sqrt{505}}{36}$, f is an almost starlike mapping of order α in D^n, and $\frac{D^2f_k(0)(z^2)}{2!}=a_kz_k^2, \frac{D^3f_k(0)(z^3)}{3!}=b_kz_k^3, k=1,2,\cdots,n, z\in D^n$, where $a_k=\frac{1}{2!}\frac{\partial^2f_k(0)}{\partial^2z_k}, b_{kl}=\frac{1}{3!}\frac{\partial^3f_k(0)}{\partial^3z_k}, k,l=1,2,\cdots,n$, then

$$\frac{\|D^4f(0)(z^4)\|}{4!}\le\frac{(1-\alpha)(3-4\alpha)(4-6\alpha)}{3}\|z\|^4, z\in D^n.$$

The above estimate is sharp.

Proof For fixed $z\in D^n\backslash\{0\}$, let $z_0=\frac{z}{\|z\|}$. Define $p_j(\xi)=\frac{g_j(\xi z_0)\|z\|}{\xi z_j}, \xi\in D$, where

$$g(z)=(g_1(z),g_2(z),\cdots,g_n(z))'=(Df(z))^{-1}f(z)$$

is a column vector in $\mathbb{C}^n$, $|z_j|=\|z\|=\max_{1\le k\le n}\{|z_k|\}$. Since $f(z)$ is an almost starlike mapping of order α in D^n, we have $p_j(\xi)\in H(D), \mathrm{Re}p_j(\xi)\ge\alpha, p_j(0)=1$, and

$$p_j(\xi)=1+\frac{D^2g_j(0)(z_0^2)}{2!}\frac{\|z\|}{z_j}\xi+\cdots+\frac{D^mg_j(0)(z_0^m)}{m!}\frac{\|z\|}{z_j}\xi^{m-1}+\cdots,\xi\in D.$$

According to Lemma 2.2, we have

$$\begin{aligned}&\Big|\frac{D^4g_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}-\frac{\frac{D^2g_j(0)(z_0^2)}{2!}\frac{\|z\|}{z_j}\frac{D^3g_j(0)(z_0^3)}{3!}\frac{\|z\|}{z_j}}{1-\alpha}+\frac{(\frac{D^2g_j(0)(z_0^2)}{2!}\frac{\|z\|}{z_j})^3}{4(1-\alpha)^2}\Big|\\ \leq\ & 2(1-\alpha)-\frac{|\frac{D^2g_j(0)(z_0^2)}{2!}\frac{\|z\|}{z_j}|^2}{2(1-\alpha)}.\end{aligned}\tag{3.12}$$

From Lemma 2.4 and the conditions of Theorem 3.4, we have

$$\begin{aligned}&\frac{D^2g_j(0)(z_0^2)}{2!}\frac{\|z\|}{z_j}=-a_j\frac{z_j}{\|z\|},\\ &\frac{D^3g_j(0)(z_0^3)}{3!}\frac{\|z\|}{z_j}=2a_j^2(\frac{z_j}{\|z\|})^2-2b_j(\frac{z_j}{\|z\|})^2,\\ &\frac{D^4g_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}=-3\frac{D^4f_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}-4a_j^3(\frac{z_j}{\|z\|})^3+7a_jb_j(\frac{z_j}{\|z\|})^3.\end{aligned}$$

Consequently, we have

$$-3\frac{D^4f_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}=\frac{D^4g_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}+4a_j^3(\frac{z_j}{\|z\|})^3-7a_jb_j(\frac{z_j}{\|z\|})^3,\tag{3.13}$$

and

$$\Big|\frac{D^4g_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}+\frac{7-8\alpha}{4(1-\alpha)^2}a_j^3(\frac{z_j}{\|z\|})^3-\frac{2}{1-\alpha}a_jb_j(\frac{z_j}{\|z\|})^3\Big|\leq 2(1-\alpha)-\frac{|a_j|^2}{2(1-\alpha)}.\tag{3.14}$$

According to Theorem 2.3, we have

$$\Big|\frac{3-4\alpha}{2(1-\alpha)}a_j^2(\frac{z_j}{\|z\|})^2-2b_j(\frac{z_j}{\|z\|})^2\Big|\leq 2(1-\alpha)-\frac{|a_j|^2}{2(1-\alpha)}.\tag{3.15}$$

Connecting (3.11), (3.12) and (3.13), we have

$$\begin{aligned}\Big|-3\frac{D^4f_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}\Big| &= \Big|\frac{D^4g_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}+4a_j^3(\frac{z_j}{\|z\|})^3-7a_jb_j(\frac{z_j}{\|z\|})^3\Big|\\ &= \Big|\Big(\frac{D^4g_j(0)(z_0^4)}{4!}\frac{\|z\|}{z_j}+\frac{7-8\alpha}{4(1-\alpha)^2}a_j^3(\frac{z_j}{\|z\|})^3-\frac{2}{1-\alpha}a_jb_j(\frac{z_j}{\|z\|})^3\Big|\Big)\\ &\quad+\frac{5-7\alpha}{2(1-\alpha)}a_j\frac{z_j}{\|z\|}\Big[\frac{3-4\alpha}{2(1-\alpha)}a_j^2(\frac{z_j}{\|z\|})^2-2b_j(\frac{z_j}{\|z\|})^2\Big]\\ &\quad-\frac{12\alpha^2-17\alpha+6}{4(1-\alpha)^2}a_j^3(\frac{z_j}{\|z\|})^3\Big|\\ &= 2(1-\alpha)-\frac{|a_j|^2}{2(1-\alpha)}+\frac{5-7\alpha}{2(1-\alpha)}|a_j|\Big(2(1-\alpha)-\frac{|a_j|^2}{2(1-\alpha)}\Big)\\ &\quad+\frac{12\alpha^2-17\alpha+6}{4(1-\alpha)^2}|a_j|^3\\ &= \frac{12\alpha^2-10\alpha+1}{4(1-\alpha)^2}|a_j|^3-\frac{|a_j|^2}{2(1-\alpha)}+(5-7\alpha)|a_j|+2(1-\alpha).\end{aligned}$$

By Lemma 3.8, we obtain

$$\begin{aligned}\Big|\frac{D^4 f_j(0)(z_0^4)}{4!}\Big| &\leqslant \frac{1}{3}\Big[\frac{12\alpha^2-10\alpha+1}{4(1-\alpha)^2}M^3-\frac{1}{2(1-\alpha)}M^2+(5-7\alpha)M|+2(1-\alpha)\Big]\\ &\leqslant \frac{(1-\alpha)(3-4\alpha)(4-6\alpha)}{3}.\end{aligned}$$

Notice that $M=\max_{\{1\leq k\leq n\}}\{|a_k|\}\leq 2(1-\alpha)$, by Lemma 3.4,

$$\frac{\|D^4 f(0)(z^4)\|}{4!}\leq\frac{(1-\alpha)(3-4\alpha)(4-6\alpha)}{3}\|z\|^4, z\in D^n.$$

Example in the proof of Theorem 3.3 verifies the accuracy of theorem 3.5. This completes the proof.

Acknowledgment

The authors thank Professor Tai-Shun Liu for providing a lot of valuable suggestions to improve the quality of this paper.

References

[1] H. Cartan, Sur la possibilité d'étendre aux fonctions de plusieurs variables complexes la theorie des fonctions univalents. In: Montel P, ed. Lecons sur les Fonctions Univalents ou Mutivalents. Paris: Gauthier-Villar, 1933. 129-155

[2] I. Graham and G. Kohr, Geometric Function Theory in One and Higher Dimensions, New York: Marcel Dekker, 2003.

[3] S. Gong, The Bieberbach Conjecture (ed. Providence, R. I.), Amer. Math. Soc., International Press, (1999) 126-176.

[4] T.J. Suffridge, Starlike and convex maps in Banach spaces, Pacific Jour. of Math., 46 (1973) 575-589.

[5] G. Kohr, On some best bounds for coefficients of several subclasses of biholomorphic mappings in Cn, Complex Variables, 36 (1998) 261-284.

[6] K.A. Roper and T.J. Suffridge, Convexity properties of holomorphic mappings in Cn, Trans.Amer.Math.Soc., 351 (1999) 1803-1833.

[7] K.A. Roper and T.J. Suffridge, Convex mappings on the unit ball of Cn, J. Anal. Math., 65 (1995) 333-347.

[8] T.S. Liu and X.S. Liu, A refinement about estimation of expansion coefficients for normalized biholomorphic mappings, Sci. China, 48 (2005) 865-879.

[9] M.S. Liu and Y.C. Zhu, The radius of convexity and the sufficient condition for starlike mappings. Bull. Malaysian Math. Sci. Soc. 35 (2012) 425-433 .

[10] X.S. Liu and T.S. Liu, The sharp estimates of all homogeneous expansions for a class of quasi-convex mappings on the unit polydisk in Cn, Chin. Ann. Math.Ser. B 32 (2012) 241-252.

[11] Y.Y. Lin and Y. Hong, Some properties of holomorphic maps in Banach spaces, Acta Math. Sinica (in Chinese), 38(2) (1995) 234-241.

[12] X.S. Liu, On the quasi-convex mappings on the unit polydisk in $\mathbb{C}^n$. J Math Anal Appl, 335 (2007) 43-55.

[13] X.S. Liu and T.S. Liu, An inequality of homogeneous expansions for biholomorphic quasi-convex mappings on the unit polydisk and its application, Acta Mathematica Scientia, 29(B)(1) (2009) 201-209.

[14] X.S. Liu and M.S. Liu, Quasi-convex mappings of order α on the unit polydisk in $\mathbb{C}^n$, Rocky Mountain J. Math., 40 (2010) 1619-1644.

[15] X.S. Liu and T.S. Liu, The sharp estimate of the third homogeneous expansions for a class of starlike mappings of order α on the unit polydisk in $\mathbb{C}^n$, Acta Mathematica Scientia, 32B(2) (2012) 752-764.

[16] Q. H. Xu and T.S. Liu, The estimation of expansion for a subclass of close-to-convex mappings, $Sci.ChinaSer.A-Math.$, 42(9) (2012) 927-938.

[17] X.S. Liu, T.S. Liu and Q.H. Xu, On sharp estimates of homogeneous expansions for starlike mappings of order α in several complex variables, Taiwanese Journal of Mathemmatics, 17(3) (2013) 801-813.

[18] X.M. Tang and M.S. Liu, Some sufficient conditions for almost spirallike mappings of order β and order α. Journal of South China Normal University, 44(3) (2012) 32-35.

[19] Q.H. Xu, T. S. Liu and X.S. Liu, The sharp estimates of all homogeneous expansions for a class of close-to-convex mappings, Annuals of Math, 33A(3)(2012) 297-310.

[20] S.Z. Deng, Y.C. Deng, C.T. Chen, C.Q. Deng and X.S. Liu, The sharp estimate of the fourth homogeneous expansions for a class of starlike mappings of order α on the unit polydisk in $\mathbb{C}^n$, Journal of Zhanjiang Normal University, 34(3) (2012) 24-30.

[21] I. Sarantopoulos, Estimates for polynomial norms on $L^p(\mu)$ spaces, Math.Proc. Cambridge Philos. Soc. 99(1986), 263-271.

[22] M.K. Papadiamantis, Polynomial estimates and radius of analyticity on real Banach spaces, arXiv:1210.7716vl [math.FA] 29 Oct 2012.